



CCPM Works Cited

1. Carr, L.J., Harrison, A.L., Evans, A.L., Stephens, J.A. Patterns of central motor reorganization in hemiplegic cerebral palsy. *Brain Oct* 1993, 116 (5) 1223-1247; DOI. 10.1093/brain/116.5.1223
2. Larson, E. B., Wang, L., Bowen, J. D., McCormick, W. C., Teri, L., Crane, P., et al. (2006). Exercise is associated with reduced risk for incident dementia among persons 65 years of age and older. *Ann. Intern. Med.* 144, 73–81. doi: 10.7326/0003-4819-144-2-200601170-00004
3. Hamer, M., and Chida, Y. (2009). Physical activity and risk of neurodegenerative disease: a systematic review of prospective evidence. *Psychol. Med.* 39, 3–11. doi: 10.1017/S0033291708003681
4. Sofi, F., Valecchi, D., Bacci, D., Abbate, R., Gensini, G. F., Casini, A., et al. (2011). Physical activity and risk of cognitive decline: a meta-analysis of prospective studies. *J. Intern. Med.* 269, 107–117. doi: 10.1111/j.1365-2796.2010.02281.x
5. Buchman, A. S., Boyle, P. A., Yu, L., Shah, R. C., Wilson, R. S., and Bennett, D. A. (2012). Total daily physical activity and the risk of AD and cognitive decline in older adults. *Neurology* 78, 1323–1329. doi: 10.1212/WNL.0b013e3182535d35
6. Xu, Q., Park, Y., Huang, X., Hollenbeck, A., Blair, A., Schatzkin, A., et al. (2010). Physical activities and future risk of Parkinson disease. *Neurology* 75, 341–348. doi: 10.1212/WNL.0b013e3181ea1597
7. Do Lee, C., Folsom, A. R., and Blair, S. N. (2003). Physical activity and stroke risk: a meta-analysis. *Stroke* 34, 2475–2481. doi: 10.1161/01.STR.0000091843.02517.9D
8. Duncan, P., Richards, L., Wallace, D., Stoker-Yates, J., Pohl, P., Luchies, C., et al. (1998). A randomized, controlled pilot study of a home-based exercise program for individuals with mild and moderate stroke. *Stroke* 29, 2055–2060. doi:10.1161/01.STR.29.10.2055
9. Gordon, N. F., Gulanick, M., Costa, F., Fletcher, G., Franklin, B. A., Roth, E. J., et al. (2004). Physical activity and exercise recommendations for stroke survivors an American heart association scientific statement from the council on clinical cardiology, subcommittee on exercise, cardiac rehabilitation, and prevention; the council on cardiovascular nursing; the council on nutrition, physical activity, and metabolism; and the stroke council. *Stroke* 35, 1230–1240. doi: 10.1161/01.STR.0000127303.19261.19
10. Crizzle, A. M., and Newhouse, I. J. (2006). Is physical exercise beneficial for persons with Parkinson's disease? *Clin. J. Sport Med.* 16, 422–425. doi:10.1097/01.jsm.0000244612.55550.7ds
11. Goodwin, V. A., Richards, S. H., Taylor, R. S., Taylor, A. H., and Campbell, J. L. (2008). The effectiveness of exercise interventions for people with Parkinson's disease: a systematic review and meta-analysis. *Mov. Disord.* 23, 631–640. doi:10.1002/mds.21922
12. Heyn, P., Abreu, B. C., and Ottenbacher, K. J. (2004). The effects of exercise training on elderly persons with cognitive impairment and dementia: a meta-analysis. *Arch. Phys. Med. Rehabil.* 85, 1694–1704. doi: 10.1016/j.apmr.2004.03.019
13. Hillman, C. H., Erickson, K. I., and Kramer, A. F. (2008). Be smart, exercise your heart: exercise effects on brain and cognition. *Nat. Rev. Neurosci.* 9, 58–65. doi:10.1038/nrn2298
14. van Praag, H. (2008). Neurogenesis and exercise: past and future directions. *Neuromolecular Med.* 10, 128–140. doi: 10.1007/s12017-008-8028-z
15. Cotman, C. W., Berchtold, N. C., and Christie, L. A. (2007). Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends Neurosci.* 30, 464–472. doi: 10.1016/j.tins.2007.06.011

16. Voss, M. W., Viviar, C., Kramer, A. F., and van Praag, H. (2013). Bridging animal and human models of exercise-induced brain plasticity. *Trends Cogn. Sci. (Regul. Ed.)* 17, 525–544. doi: 10.1016/j.tics.2013.08.001
17. Kramer, A. F., and Erickson, K. I. (2007). Capitalizing on cortical plasticity: influence of physical activity on cognition and brain function. *Trends Cogn. Sci. (Regul. Ed.)* 11, 342–348. doi: 10.1016/j.tics.2007.06.009
18. Kleim, J. A., Cooper, N. R., and VandenBerg, P. M. (2002). Exercise induces angio-genesis but does not alter movement representations within rat motor cortex. *Brain Res.* 934, 1–6. doi: 10.1016/S0006-8993(02)02239-4
19. Swain, R. A., Harris, A. B., Wiener, E. C., Dutka, M. V., Morris, H. D., Theien, B. E., et al. (2003). Prolonged exercise induces angiogenesis and increases cerebral blood volume in primary motor cortex of the rat. *Neuroscience* 117, 1037–1046. doi: 10.1016/S0306-4522(02)00664-4
20. Cotman, C. W., and Berchtold, N. C. (2002). Exercise: a behavioral intervention to enhance brain health and plasticity. *Trends Neurosci.* 25, 295–301. doi:10.1016/S0166-2236(02)02143-4
21. Vaynman, S., and Gomez-Pinilla, F. (2005). License to run: exercise impacts functional plasticity in the intact and injured central nervous system by using neurotrophins. *Neurorehabil. Neural Rep.* 19, 283–295. doi:10.1177/1545968305280753
22. Paltsev YI, Elnor AM. Change in the functional state of the segmental apparatus of the spinal cord under the influence of sound stimuli and its role in voluntary movement. *Biophysics.* 1967;12:1219–26.
23. Rossignol S, Melvill Jones G. Audiospinal influences in man studied by the H-reflex and its possible role in rhythmic movement synchronized to sound. *Electroencephalogr Clin Neurophysiol.* 1976;41:83–92.
24. Thaut, M. H., and Abiru, M. (2010). Rhythmic auditory stimulation in rehabilitation of movement disorders: a review of current research. *Music Percept.* 27, 263–269 doi: 10.1525/mp.2010.27.4.263
25. Hurt, C. P., Rice, R. R., McIntosh, G. C., and Thaut, M. H. (1998). Rhythmic auditory stimulation in gait training for patients with traumatic brain injury. *J. Music Ther.* 35, 228–241. doi: 10.1093/jmt/35.4.228
26. Thaut, M. H., Gardiner, J. C., Holmberg, D., Horwitz, J., Kent, L., Andrews, G., et al. (2009). Neurologic music therapy improves executive function and emotional adjustment in traumatic brain injury rehabilitation. *Ann. N. Y. Acad. Sci.* 1169, 406–416. doi: 10.1111/j.1749-6632.2009.04585.x
27. Schlaug, G., Norton, A., Marchina, S., Zipse, L., and Wan, C. Y. (2010). From singing to speaking: facilitating recovery from nonfluent aphasia. *Future Neurol.* 5, 657–665. doi: 10.2217/fnl.10.44
28. Thaut MH, McIntosh GC, Prassas SG, Rice RR. The effect of auditory rhythmic cuing on stride and EMG patterns in hemiparetic gait of stroke patients. *J Neurol Rehabil.* 1993;7:9–16.
29. Grau-Sánchez J, Amengual JL, Rojo N, et al. Plasticity in the sensorimotor cortex induced by Music-supported therapy in stroke patients: a TMS study. *Frontiers in Human Neuroscience.* 2013;7:494. doi:10.3389/fnhum.2013.00494
30. Johnson, G., Otto, D., and Clair, A. A. (2001). The effect of instrumental and vocal music on adherence to a physical rehabilitation exercise program with persons who are elderly. *J. Music Ther.* 38, 82–96. doi: 10.1093/jmt/38.2.82
31. Schutzer, K. A., and Graves, B. S. (2004). Barriers and motivations to exercise in older adults. *Prev. Med.* 39, 1056–1061. doi: 10.1016/j.ypmed.2004.04.003
32. Kattenstroth, J. C., Kolankowska, I., Kalisch, T., and Dinse, H. R. (2010). Superior sensory, motor, and cognitive performance in elderly individuals with multi-year dancing activities. *Front. Aging Neurosci.* 2:31. doi: 10.3389/fnagi.2010.00031
33. Fissler, P., Küster, O., Schlee, W., and Kolassa, I. T. (2012). Novelty interventions to enhance broad cognitive abilities and prevent dementia: synergistic approaches for the facilitation of positive plastic change. *Prog. Brain Res.* 207, 403–434. doi: 10.1016/B978-0-444-63327-9.00017-5
34. Kraft, E. (2012). Cognitive function, physical activity, and aging: possible biological links and implications for multimodal interventions. *Aging Neuropsychol. Cogn.* 19, 248–263. doi: 10.1080/13825585.2011.645010
35. Olson, A. K., Eadie, B. D., Ernst, C., and Christie, B. R. (2006). Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. *Hippocampus* 16, 250–260. doi:10.1002/hipo.20157
36. Foster, P. P. (2013). How does dancing promote brain reconditioning in the elderly? *Front. Aging Neurosci.* 5:4. doi: 10.3389/fnagi.2013.00004
37. Johansson, B. B. (2012). Multisensory stimulation in stroke rehabilitation. *Front. Hum. Neurosci.* 6:60. doi: 10.3389/fnhum.2012.00060
38. Hänggi, J., Koeneke, S., Bezzola, L., and Jäncke, L. (2010). Structural neuroplasticity in the sensorimotor network of professional female ballet dancers. *Hum. Brain Mapp.* 31, 1196–1206.
39. Hüfner, K., Binetti, C., Hamilton, D. A., Stephan, T., Flanagin, V. L., Linn, J., et al. (2011). Structural and functional plasticity of the hippocampal formation in professional dancers and slackliners. *Hippocampus* 21, 855–865. doi:10.1002/hipo.20801
40. Kempermann, G., Fabel, K., Ehninger, D., Babu, H., Leal-Galicia, P., Garthe, A., et al. (2010). Why and how physical

- activity promotes experience-induced brain plasticity. *Front. Neurosci.* 4:189. doi: 10.3389/fnins.2010.00189
41. Fabel, K., Wolf, S. A., Ehninger, D., Babu, H., Leal-Galicia, P., and Kempermann, G. (2009). Additive effects of physical exercise and environmental enrichment on adult hippocampal neurogenesis in mice. *Front. Neurosci.* 3:50. doi: 10.3389/neuro.22.002.2009
 42. Fabel, K., and Kempermann, G. (2008). Physical activity and the regulation of neurogenesis in the adult and aging brain. *Neuromolecular Med.* 10, 59–66. doi:10.1007/s12017-008-8031-4
 43. Curlik, D. M. II, Maeng, L. Y., Agarwal, P. R., and Shors, T. J. (2011). Gym Rats are Not as Stupid as You Think: Physical Skill Learning Increases Cell Survival in the Hippocampus. Program No. 595.10. Washington, DC: Neuroscience Meeting Planner, Society for Neuroscience.
 44. Hopkins, D. R., Murrah, B., Hoeger, W. W., and Rhodes, R. C. (1990). Effect of low-impact aerobic dance on the functional fitness of elderly women. *Gerontologist* 30, 189–192. doi: 10.1093/geront/30.2.189
 45. Kattenstroth, J. C., Kalisch, T., Holt, S., Tegenthoff, M., and Dinse, H. R. (2013). Six months of dance intervention enhances postural, sensorimotor, and cognitive performance in elderly without affecting cardio-respiratory functions. *Front. Aging Neurosci.* 5:5. doi: 10.3389/fnagi.2013.00005
 46. Keogh, J. W., Kilding, A., Pidgeon, P., Ashley, L., and Gillis, D. (2009). Physical benefits of dancing for healthy older adults: a review. *J. Aging Phys. Act.* 17, 479–500.
 47. Verghese, J. (2006). Cognitive and mobility profile of older social dancers. *J. Am. Geriatr. Soc.* 54, 1241–1244. doi: 10.1111/j.1532-5415.2006.00808.x
 48. Federici, A., Bellagamba, S., and Rocchi, M. B. (2005). Does dance-based training improve balance in adult and young old subjects? A pilot randomized controlled trial. *Aging Clin. Exp. Res.* 17, 385–389. doi: 10.1007/BF03324627
 49. Hui, E., Chui, B. T. K., and Woo, J. (2009). Effects of dance on physical and psychological well-being in older persons. *Arch. Gerontol. Geriatr.* 49, e45–e50. doi: 10.1016/j.archger.2008.08.006
 50. Shigematsu, R., Chang, M., Yabushita, N., Sakai, T., Nakagaichi, M., Nho, H., et al. (2002). Dance-based aerobic exercise may improve indices of falling risk in older women. *Age Ageing* 31, 261–266. doi: 10.1093/ageing/31.4.261
 51. Duncan, R. P., and Earhart, G. M. (2012). Randomized controlled trial of community-based dancing to modify disease progression in Parkinson disease. *Neurorehabil. Neural Rep.* 26, 132–143. doi: 10.1177/1545968311421614
 52. Earhart, G. M. (2009). Dance as therapy for individuals with Parkinson disease. *Eur. J. Phys. Rehabil. Med.* 45, 231–238.
 53. Hackney, M. E., and Earhart, G. M. (2010). Effects of dance on balance and gait in severe Parkinson disease: a case study. *Disabil. Rehabil.* 32, 679–684. doi:10.3109/09638280903247905
 54. Hackney, M. E., Hall, C. D., Echt, K.V., and Wolf, S. L. (2012). Application of adapted tango as therapeutic intervention for patients with chronic stroke. *J. Geriatr. Phys. Ther.* 35, 206–217. doi: 10.1519/JPT.0b013e31823ae6ea
 55. Heiberger, L., Maurer, C., Amtage, F., Mendez-Balbuena, I., Schulte-Mönting, J., Hepp-Reymond, M. C., et al. (2011). Impact of a weekly dance class on the functional mobility and on the quality of life of individuals with Parkinson's disease. *Front. Aging Neurosci.* 3:14. doi: 10.3389/fnagi.2011.00014
 56. Stranahan, A. M., Khalil, D., and Gould, E. (2006). Social isolation delays the positive effects of running on adult neurogenesis. *Nat. Neurosci.* 9, 526–533. doi:10.1038/nn1668
 57. Saarela, M. V., and Hari, R. (2008). Listening to humans walking together activates the social brain circuitry. *Soc. Neurosci.* 3, 401–409. doi:10.1080/17470910801897633
 58. Van Der Bij, A. K., Laurant, M. G., and Wensing, M. (2002). Effectiveness of physical activity interventions for older adults: a review. *Am. J. Prev. Med.* 22, 120–133. doi: 10.1016/S0749-3797(01)00413-5
 59. Wankel, L. M. (1985). Personal and situational factors affecting exercise involvement: the importance of enjoyment. *Res. Q. Exerc. Sport* 56, 275–282. doi:10.1080/02701367.1985.10605374
 60. Dishman, R. K. (1988). *Exercise Adherence: Its Impact on Public Health*. Champaign: Human Kinetics Publishers
 61. Rhodes, R. E., Martin, A. D., Taunton, J. E., Rhodes, E. C., Donnelly, M., and Elliot, J. (1999). Factors associated with exercise adherence among older adults. *Sports Med.* 28, 397–411. doi: 10.2165/00007256-199928060-00003